

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : B01D 15/08	A1	(11) International Publication Number: WO 95/22389 (43) International Publication Date: 24 August 1995 (24.08.95)
(21) International Application Number: PCT/US95/01823 (22) International Filing Date: 21 February 1995 (21.02.95) (30) Priority Data: 08/200,126 22 February 1994 (22.02.94) US (60) Parent Application or Grant (63) Related by Continuation US 08/200,126 (CON) Filed on 22 February 1994 (22.02.94) (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SHADLE, Paula, J. [US/US]; 425 Garrison Way, Gulph Mills, PA 19428 (US). ERICKSON, John, C. [US/US]; 201 Cardinal Drive, Conshohocken, PA 19428 (US). SCOTT, Robert, G. [US/US]; 827 Stokes Avenue, Collingswood, NJ 08108 (US). SMITH, Thomas, M. [US/US]; 4009 Marshall Road, Drexel Hill, PA 19026 (US).	(74) Agents: JERVIS, Herbert, H. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US). (81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, MX, NO, NZ, PT, RO, RU, SD, SI, SK, UA, US, VN, European patent (BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: ANTIBODY PURIFICATION (57) Abstract This invention relates to the application of hydrophobic interaction chromatography combination chromatography to the purification of antibody molecule proteins. The process may include the sequential steps of Protein A affinity chromatography, ion exchange chromatography, and hydrophobic interaction chromatography. Purified monomeric IgG antibody, free of immunoglobulin aggregates, misfolded species, host cell protein, and Protein A, may be recovered by the process.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Antibody Purification

Field of the Invention:

This invention relates to the field of protein purification. More specifically, this invention relates to the application of Hydrophobic Interaction Chromatography (HIC) to the separation of Immunoglobulin G monomers and to the integration of HIC into a combination chromatographic protocol for the purification of IgG antibody molecules.

Background of the Invention:

Historically, protein purification schemes have been predicated on differences in the molecular properties of size, charge and solubility between the protein to be purified and undesired protein contaminants. Protocols based on these parameters include size exclusion chromatography, ion exchange chromatography, differential precipitation and the like.

Size exclusion chromatography, otherwise known as gel filtration or gel permeation chromatography, relies on the penetration of macromolecules in a mobile phase into the pores of stationary phase particles. Differential penetration is a function of the hydrodynamic volume of the particles. Accordingly, under ideal conditions the larger molecules are excluded from the interior of the particles while the smaller molecules are accessible to this volume and the order of elution can be predicted by the size of the protein because a linear relationship exists between elution volume and the log of the molecular weight.

Chromatographic supports based on cross-linked dextrans e.g. SEPHADEX[®], spherical agarose beads e.g. SEPHAROSE[®] (both commercially available from Pharmacia AB, Uppsala, Sweden), based on cross-linked polyacrylamides e.g. BIO-GEL[®] (commercially available from BioRad Laboratories, Richmond, California) or based on ethylene glycol-methacrylate copolymer e.g. TOYOPEARL HW65 (commercially available from Toso Haas Co., Tokyo, Japan) are useful in forming the various chromatographic columns for size exclusion, or HIC chromatography in the practice of certain aspects of this invention.

Precipitation methods are predicated on the fact that in crude mixtures of proteins the solubilities of individual proteins are likely to vary widely. Although the solubility of a protein in an aqueous medium depends on a variety of factors, for purposes of this discussion it can be said generally that a protein

will be soluble if its interaction with the solvent is stronger than its interaction with protein molecules of the same or similar kind. Without wishing to be bound by any particular mechanistic theory describing precipitation phenomena, it is nonetheless believed that interaction between a protein and water molecules can occur by hydrogen bonding with several types of uncharged groups and/or electrostatically, as dipoles, with charged groups and that precipitants such as salts of monovalent cations (e.g. ammonium sulfate) compete with proteins for water molecules. Thus at high salt concentrations, the proteins become "dehydrated" reducing their interaction with the aqueous environment and increasing the aggregation with like or similar proteins, resulting in precipitation from the medium.

Ion exchange chromatography involves the interaction of charged functional groups in the sample with ionic functional groups of opposite charge on an adsorbent surface. Two general types of interaction are known. Anionic exchange chromatography is mediated by negatively charged amino acid side chains (e.g., aspartic acid and glutamic acid) interacting with positively charged surfaces and cationic exchange chromatography is mediated by positively charged amino acid residues (e.g., lysine and arginine) interacting with negatively charged surfaces.

More recently affinity chromatography and hydrophobic interaction chromatography techniques have been developed to supplement the more traditional size exclusion and ion exchange chromatographic protocols. Affinity chromatography relies on the specific interaction of the protein with an immobilized ligand. The ligand can be specific for the particular protein of interest in which case the ligand is a substrate, substrate analog, inhibitor, receptor or antibody. Alternatively, the ligand may be able to react with a number of related proteins. Such group specific ligands as adenosine monophosphate, adenosine diphosphate, nicotine adenine dinucleotide or certain dyes may be employed to recover a particular class of proteins.

With respect to the purification of antibody molecules, both specific and generalized affinity techniques are applicable. The most specific choice of ligand for the affinity purification of an antibody is the antigen (or an epitope thereof) to which desired antibody reacts. Many of the well-known immunosorbent assays such as the enzyme-linked immunosorbent assays (ELISA) are predicated on such specific antigen/antibody affinity interactions.

However, generalized affinity techniques are also useful. For example, Staphylococcal Protein A is known to bind certain antibodies of the IgG class

(See: Ey, P.L. *et al.* Immunochimistry 15:429-36 (1978)). Alternatively, antisera raised in heterologous species (e.g. rabbit anti-mouse antisera) can be used to separate general groups of antibodies. (See, Current Protocols in Molecular Biology *Supra*, Chap 11.)

5 Hydrophobic interaction chromatography was first developed following the observation that proteins could be retained on affinity gels which comprised hydrocarbon spacer arms but lacked the affinity ligand. Although in this field the term hydrophobic chromatography is sometimes used, the term hydrophobic interaction chromatography (HIC) is preferred because it is the interaction
10 between the solute and the gel that is hydrophobic not the chromatographic procedure. Hydrophobic interactions are strongest at high ionic strength, therefore, this form of separation is conveniently performed following salt precipitations or ion exchange procedures. Elution from HIC supports can be effected by alterations in solvent, pH, ionic strength, or by the addition of
15 chaotropic agents or organic modifiers, such as ethylene or propylene glycol. A description of the general principles of hydrophobic interaction chromatography can be found in U. S. Patent 3,917,527 and in U. S. Patent 4,000,098. The application of HIC to the purification of specific proteins is exemplified by reference to the following disclosures: human growth hormone (U. S. Patent
20 4,332,717), toxin conjugates (U. S. Patent 4,771,128), antihemolytic factor (U. S. Patent 4,743,680), tumor necrosis factor (U. S. Patent 4,894,439), interleukin-2 (U. S. Patent 4,908,434), human lymphotoxin (U. S. Patent 4,920,196) and lysozyme species (Fausnaugh, J.L. and F. E. Regnier, J. Chromatog. 359:131-146 (1986)) and soluble complement receptors (U.S. Patent 5,252,216). HIC in
25 the context of high performance liquid chromatography (HPLC) has been used to separate antibody fragmens (e.g., F(ab')₂) from intact antibody molecules in a single step protocol. (Morimoto, K. *et al.*, J. Biochem. Biophys. Meth. 24: 107-117 (1992)).

 In addition to affinity and HIC techniques, one or more of the traditional
30 protein purification schemes have been applied to antibody purification. For example, Hakalahti, L. *et al.*, (J. Immunol. Meth. 117: 131-136 (1989)) disclose a protocol employing two successive ion exchange chromatographic steps or one employing a single ion exchange step followed by a HIC step. Danielsson A. *et al.* (J. Immunol. Methods 115: 79-88 (1988)) compare single step protocols
35 based on anion exchange, cation exchange, chromatofocusing and HIC respectively.

Although Protein A affinity column chromatography is widely used, it is also appreciated that elution of antibody from such columns can result in leaching of residual Protein A from the support. Size exclusion HPLC (Das *et al.*, Analytical Biochem. 145: 27-36 (1985)) and anion exchange chromatography (EPO345549, published Dec. 13, 1989) have been suggested as means for dealing with this problem.

It has now been surprisingly discovered that HIC can be usefully employed to remove contaminating Protein A from IgG mixtures eluted from Protein A chromatographic support.

This invention relates to the application of HIC to the separation of monomeric IgG from mixtures containing same and to the integration of HIC into a protocol combining Protein A and ion exchange chromatography for the purification of immunoglobulin G molecules.

Brief Description of the Invention:

This invention relates to a method for separating IgG monomers from aggregates in mixtures containing same by contacting said mixture with a hydrophobic interaction chromatographic support and selectively eluting the monomer from the support.

In another aspect the invention provides for the purification of an IgG antibody from conditioned cell culture medium containing same comprising sequentially subjecting the medium to (a) Protein A, (b) ion exchange chromatography, and (c) hydrophobic interaction chromatography.

In another aspect the invention provides a method for removing Protein A from a mixture comprising Protein A and antibodies comprising contacting said mixture with a hydrophobic interaction chromatography support and selectively eluting the antibody from the support.

Detailed Description of the Figures:

Figure 1 illustrates a flow diagram of one process for purifying an antibody according to this invention.

Detailed Description of the Invention:

This invention relates to protein purification techniques which have application to the large scale purification of immunoglobulin molecules. The invention is particularly useful because it permits the recovery of monomeric

IgG of >95% protein purity. The invention may be applied to the purification of a number of different immunoglobulin G molecules.

Antibody-like proteins are proteins which may be purified by the protocol described herein, such protocol being modified if necessary by routine, non-inventive adjustments that do not entail undue experimentation. Such proteins include isotypes, allotypes and alleles of immunoglobulin genes, truncated forms, altered antibodies, such as chimeric antibodies, humanized antibodies and the like, chemically modified forms such as by PEG treatment, and fusion proteins containing an immunoglobulin moiety. These proteins are referred to as antibody-like because they possess or retain sufficient immunoglobulin protein properties (e.g. F_C determinants) to admit to purification by the process of this invention. Unless specifically identified otherwise, the term antibody or immunoglobulin protein also includes antibody-like proteins.

The immunoglobulin molecules of this invention can be isolated from a number of sources, including without limitation, serum of immunized animals, ascites fluid, hybridoma or myeloma supernatants, conditioned media derived from culturing a recombinant cell line that expresses the immunoglobulin molecule and from all cell extracts of immunoglobulin producing cells. This invention is particularly useful for the purification of antibodies from conditioned cell culture media of a variety of antibody producing recombinant cell lines. Although one may expect some variation from cell line to cell line and among the various antibody products, based on the disclosure herein, it is well within the purview of one of ordinary skill in this art to adapt the invention herein to a particular combination of antibody protein and producing cell line.

Generally, genes encoding proteins such as antibodies may be cloned by incorporating DNA sequences coding for the desired regions of the polypeptide into a recombinant DNA vehicle (e.g., vector) and transforming or transfecting suitable prokaryotic or eukaryotic hosts. Suitable prokaryotic hosts include but are not limited to Escherichia, Streptomyces, Bacillus and the like. Suitable eukaryotic hosts include but are not limited to yeast, such as Saccharomyces and animal cells in culture such as VERO, HeLa, mouse C127, Chinese hamster ovary (CHO), WI-38, BHK, COS, MDCK, myeloma, and insect cell lines. Particularly preferred hosts are CHO cell lines deficient in dihydrofolate reductase such as ATCC CRL 1793, CRL 9096 and other cell lines described herein below. Such recombinant techniques have now become well known and are described in Methods in Enzymology, (Academic Press) Volumes 65 and 69

(1979), 100 and 101 (1983), and the references cited therein. An extensive technical discussion embodying most commonly used recombinant DNA methodologies can be found in Maniatis, *et al.*, Molecular Cloning, Cold Spring Harbor Laboratory (1982) or Current Protocols in Molecular Biology, Greene Publishing, Wiley Interscience (1988,1991,1993).

One way of obtaining a DNA fragment encoding a desired polypeptide such as an antibody molecule is via cDNA cloning. In this process, messenger RNA (mRNA) is isolated from cells known or suspected of producing the desired protein. Through a series of enzymatic reactions, the mRNA population of the cells is copied into a complementary DNA (cDNA). The resulting cDNA is then inserted into cloning vehicles and subsequently used to transform a suitable prokaryotic or eukaryotic host. The resulting cDNA "library" is comprised of a population of transformed host cells, each of which contain a single gene or gene fragment. The entire library, in theory, provides a representative sample of the coding information present in the mRNA mixture used as the starting material. The libraries can be screened using nucleic acid or antibody probes in order to identify specific DNA sequences. Once isolated, these DNA sequences can be modified or can be assembled into complete genes.

Specific fragments of an antibody gene can be engineered independently of the rest of the gene. DNA fragments encoding Complementarity Determining Regions (CDRs) can be integrated into DNA framework sequences from heterologous species to yield altered antibodies. These altered antibodies have significant utility in the treatment of undesirable physiological conditions. For example, PCT/GB91/01554 (published as WO92/04381) discloses the production of "humanized" antibodies useful for the treatment and prevention of Respiratory Syncytial Virus (RSV) infection. Alternatively, the entire variable region of antibody gene can be fused to the constant domain of a second antibody to form an altered antibody otherwise known as a "chimeric antibody". For example, PCT/US92/06194 (published as WO93/02108) discloses a monkey/human chimeric antibody reactive with the human CD4 receptor.

Once the antibody gene or gene fragment has been cloned, the DNA may be introduced into an expression vector and that construction used to transform an appropriate host cell. An expression vector is characterized as having expression control sequences as defined herein, such that when a DNA sequence of interest is operably linked thereto, the vector is capable of directing the production of the product encoded by the DNA sequence of interest in a host cell containing the vector. With specific reference to this invention, it is

possible to assemble fragments of a single coding sequence such that upon expression an antibody molecule is formed. A particularly efficacious application of this protocol to recombinant antibody production is found in the Harris, et al. PCT Applications WO92/04381, published March 19, 1992, cited
5 above, and in the Newman et al. PCT Application WO93/02108, published February 4, 1993, cited above.

After the recombinant product is produced it is desirable to recover the product. If the product is exported by the cell producing it, the product can be recovered directly from the cell culture medium. If the product is retained
10 intracellularly, the cells must be physically disrupted by mechanical, chemical or biological means in order to obtain the intracellular product.

In the case of a protein product, the purification protocol should not only provide a protein product that is essentially free of other proteins, by which is meant at least 80% and preferably greater than 95% pure with respect to total
15 protein in the preparation, but also eliminate or reduce to acceptable levels other host cell contaminants, DNA, RNA, potential pyrogens and the like. Furthermore, in the context of antibody production by recombinant expression system, it is appreciated that aggregation of the 150,000 dalton IgG product into higher molecular weight species can occur. Accordingly, for purposes of
20 product purity and standardization it is also useful to separate the native 150,000 dalton monomeric species from higher molecular weight aggregates and other misfolded forms. While it is appreciated that the 150,000 dalton IgG species is composed of four polypeptide chains (2 heavy chains and 2 light chains), the 150,000 dalton species is referred to herein as a "monomer" or "monomeric
25 IgG".

As mentioned above, a variety of host cells may be used for the production of the antibodies of this invention. The choice of a particular host cell is well within the purview of the ordinary skilled artisan taking into account, inter alia, the nature of the antibody, its rate of synthesis, its rate of decay and
30 the characteristics of the recombinant vector directing the expression of the antibody. The choice of the host cell expression system dictates to a large extent the nature of the cell culture procedures to be employed. The selection of a particular mode of production, be it batch or continuous, spinner or air lift, liquid or immobilized can be made once the expression system has been
35 selected. Accordingly, fluidized bed bioreactors, hollow fiber bioreactors, roller bottle cultures, or stirred tank bioreactors, with or without cell microcarriers may variously be employed. The criteria for such selection are appreciated in

the cell culture art. They are not detailed herein because they are outside the scope of this invention. This invention relates to the purification of antibodies given their existence in a conditioned cell culture medium, hybridoma supernatant, antiserum, myeloma supernatant or ascites fluid.

5 As mentioned above this invention relates, inter alia, to application of hydrophobic interaction chromatography (HIC) to the separation and purification of antibody molecules. Hydrophobic molecules in an aqueous solvent will self-associate. This association is due to hydrophobic interactions. It is now appreciated that macromolecules such as proteins have on their surface
10 extensive hydrophobic patches in addition to the expected hydrophilic groups. HIC is predicated, in part, on the interaction of these patches with hydrophobic ligands attached to chromatographic supports. A hydrophobic ligand coupled to a matrix is variously referred to herein as an HIC support, HIC gel or HIC
15 column. It is further appreciated that the strength of the interaction between the protein and the HIC support is not only a function of the proportion of non-polar to polar surfaces on the protein but by the distribution of the non-polar surfaces as well and the chemistry of the HIC support.

A number of chromatographic supports may be employed in the preparation of HIC columns, the most extensively used are agarose, silica and
20 organic polymer or co-polymer resins. Useful hydrophobic ligands include but are not limited to alkyl groups having from about 2 to about 8 carbon atoms, such as a butyl, propyl, or octyl; or aryl groups such as phenyl. Conventional HIC products for gels and columns may be obtained commercially from suppliers such as Pharmacia LKB AB, Uppsala, Sweden under the product
25 names butyl-SEPHAROSE®, phenyl or butyl-SEPHAROSE® CL-4B, butyl-SEPHAROSE® FF, octyl-SEPHAROSE® FF and phenyl-SEPHAROSE® FF; Tosoh Corporation, Tokyo, Japan under the product names TOYOPEARL ether 650, phenyl 650 or butyl 650 (Fractogel); Miles-Yeda, Rehovot, Israel under the product name alkyl-agarose, wherein the alkyl group contains from 2-10 carbon
30 atoms, and J.T. Baker, Phillipsburg, N.J. under the product name Bakerbond WP-HI-propyl.

It is also possible to prepare the desired HIC column using conventional chemistry. (See: for example, Er-el, Z. et al. Biochem. Biophys. Res. Comm. 49:383 (1972) or Ulbrich, V. et al. Coll. Czech. Chem. Commun. 9:1466
35 (1964)).

Ligand density is an important parameter in that it influences not only the strength of the interaction but the capacity of the column as well. The ligand

density of the commercially available phenyl or octyl phenyl gels is on the order of 40 μ moles/ml gel bed. Gel capacity is a function of the particular protein in question as well as pH, temperature and salt type and concentration but generally can be expected to fall in the range of 3-20 mg/ml of gel.

5 The choice of a particular gel can be determined by the skilled artisan. In general the strength of the interaction of the protein and the HIC ligand increases with the chain length of the alkyl ligands but ligands having from about 4 to about 8 carbon atoms are suitable for most separations. A phenyl group has about the same hydrophobicity as a pentyl group, although the
10 selectivity can be quite different owing to the possibility of pi-pi orbital interaction with aromatic groups on the protein. Selectivity may also be affected by the chemistry of the supporting resin.

 Adsorption of the proteins to a HIC column is favored by high salt concentrations, but the actual concentrations can vary over a wide range
15 depending on the nature of the protein and the particular HIC ligand chosen. Various ions can be arranged in a so-called soluphobic series depending on whether they promote hydrophobic interactions (salting-out effects) or disrupt the structure of water (chaotropic effect) and lead to the weakening of the hydrophobic interaction. Cations are ranked in terms of increasing salting out
20 effect as $\text{Ba}^{++} < \text{Ca}^{++} < \text{Mg}^{++} < \text{Li}^+ < \text{Cs}^+ < \text{Na}^+ < \text{K}^+ < \text{Rb}^+ < \text{NH}_4^+$, while anions may be ranked in terms of increasing chaotropic effect as $\text{PO}_4^{---} < \text{SO}_4^{--} < \text{CH}_3\text{COO}^- < \text{Cl}^- < \text{Br}^- < \text{NO}_3^- < \text{ClO}_4^- < \text{I}^- < \text{SCN}^-$.

Accordingly, salts may be formulated that influence the strength of the interaction as given by the following relationship:

25 $(\text{NH}_4)_2\text{SO}_4 > \text{Na}_2\text{SO}_4 > \text{NaCl} > \text{NH}_4\text{Cl} > \text{NaBr} > \text{NaSCN}$

In general, salt concentrations of between about 0.75 and about 2M ammonium sulfate or between about 1 and 4M NaCl are useful.

 The influence of temperature on HIC separations is not simple, although generally a decrease in temperature decreases the interaction. However, any
30 benefit that would accrue by increasing the temperature must also be weighed against adverse effects such an increase may have on the stability of the protein.

 Elution, whether stepwise or in the form of a gradient, can be accomplished in a variety of ways: (a) by changing the salt concentration, (b) by changing the polarity of the solvent or (c) by adding detergents. By decreasing
35 salt concentration adsorbed proteins are eluted in order of increasing hydrophobicity. Changes in polarity may be affected by additions of solvents such as ethylene or propylene glycol or (iso)propanol, thereby decreasing the

strength of the hydrophobic interactions. Detergents function as displacers of proteins and have been used primarily in connection with the purification of membrane proteins.

Although it has been discovered that HIC chromatography can be used
5 alone to separate monomeric IgG (MW 150,000) from aggregates and misfolded species, as mentioned above, HIC is particularly useful when used in combination with other protein purification techniques. That is to say, it is preferred to apply HIC to mixtures that have been partially purified by other protein purification procedures. By the term "partially purified" is meant a
10 protein preparation in which the protein of interest is present in at least 5 percent by weight, more preferably at least 10% and most preferably at least 45%. By the term "mixture" is meant the desired monomeric IgG antibody molecule in combination with undesirable contaminants such as, without limitation, one or more of: immunoglobulin aggregates, misfolded species, host cell protein,
15 residue material from preceding chromatographic steps such as Protein A when employed. Accordingly, the application of HIC can also be appreciated in the context of an overall purification protocol for immunoglobulin proteins such as affinity purified monoclonal antibodies. It has been found to be useful, for example, to subject a sample of conditioned cell culture medium to partial
20 purification prior to the application of HIC. By the term "conditioned cell culture medium" is meant a cell culture medium which has supported cell growth and/or cell maintenance and contains secreted product. A sample of such medium is subjected to one or more protein purification steps prior to the application of a HIC step. The sample may be subjected to affinity
25 chromatography employing Staphylococcus Protein A as a first step. For example, PROSEP-A[®] (BioProcessing Ltd., U.K.) which consists of Protein A covalently coupled to controlled pore glass can be usefully employed. Other useful Protein A formulations are Protein A SEPHAROSE[®] Fast Flow (Pharmacia) and TOYOPEARL 650M Protein A (TosoHaas). As a second step,
30 ion exchange chromatography may be employed. In this regard various anionic or cationic substituents may be attached to matrices in order to form anionic or cationic supports for chromatography. Anionic exchange substituents include diethylaminoethyl(DEAE), quaternary aminoethyl(QAE) and quaternary amine(Q) groups. Cationic exchange substituents include carboxymethyl (CM),
35 sulfoethyl(SE), sulfopropyl(SP), phosphate(P) and sulfonate(S). Cellulosic ion exchange resins such as DE23, DE32, DE52, CM-23, CM-32 and CM-52 are available from Whatman Ltd. Maidstone, Kent, U.K. SEPHADEX[®]-based and

cross-linked ion exchangers are also known. For example, DEAE-, QAE-, CM-, and SP- SEPHADEX® and DEAE-, Q-, CM- and S-SEPHAROSE® and SEPHAROSE® Fast Flow are all available from Pharmacia AB. Further, both DEAE and CM derivitized ethylene glycol-methacrylate copolymer such as
5 TOYOPEARL DEAE-650S or M and TOYOPEARL CM-650S or M are available from Toso Haas Co., Philadelphia, Pa. Because elution from ion exchange supports usually involves addition of salt and because, as mentioned previously, HIC is enhanced under increased salt concentrations, the introduction of a HIC step following an ionic exchange chromatographic step or
10 other salt mediated purification step is particularly preferred. Additional purification protocols may be added including but not necessarily limited to further ionic exchange chromatography, size exclusion chromatography, viral inactivation, concentration and freeze drying.

For purposes of illustration only, this invention was applied to the
15 purification of several antibodies of the IgG isotype. More specifically, to a humanized antibody useful for the treatment of RSV infection described by Harris *et al.*; 1992, Intl. Patent Publication Number WO92/04381, published March 19, 1992 (hereinafter "RSHZ-19") and a chimeric antibody specifically reactive with the CD4 antigen described by Newman *et al.* Intl Patent
20 Publication Number WO93/02108, published February 4, 1993 (hereinafter CH-CD4). The construction of recombinant systems for the production of RSHZ-19 and the CH-CD4 chimeric antibodies are detailed in the above mentioned PCT Applications, the contents of which are incorporated herein by reference for purpose of background and are summarized as follows.

25 An expression plasmid containing the RSHZ-19 coding sequence was co-transfected with pSV2dhfr into a dhfr-requiring Chinese Hamster Ovary cell line (CHODUXBII). The transfection was carried in growth medium and employed the calcium coprecipitation/ glycerol shock procedure as described in: DNA Cloning, D.M. Glover ed. (Chap. 15, C. Gorman). Following transfection, the
30 cells were maintained in growth medium for 46 hours under growth conditions (as described above) prior to the selection procedure.

The selection and co-amplification procedure was carried out essentially as described by R.J. Kaufman, *et al.* (*Mol. Cell. Biol.* 5:1750-1759 (1985)). Forty-six hours post transfection the cells were changed to selective medium
35 MEM ALPHA (041-02571), 1% stock glutamine, 1% stock pen/strep (043-05070) and dialyzed bovine fetal calf serum (220-6300AJ) (Gibco, Paisley, Scotland). The cells were maintained in the selective medium for 8-10 days

until dhfr⁺ colonies appeared. When the colonies were established the cells were changed into a selective medium containing methotrexate, (A6770, Sigma Chem. Co., St. Louis, Mo.). The methotrexate concentration was initially 0.02µM and was increased stepwise to 5µM. During the amplification procedure
5 aliquots of growth medium from growing cells were assayed for RSHZ-19 production by human IgG. Any antibody secreting recombinant cell line may be used to supply the conditioned medium for purification according to this invention, a particular cell line certainly is not required.

10 A transfected CHO cell line capable of producing RSHZ-19 can be cultured by a variety of cell culture techniques. For the application of this invention the particular method of culturing is not critical.

As mentioned previously, the particular recombinant production system and the particular cell culturing protocol is outside the scope of this invention. The system and protocol discussed above are representative of the many options
15 available to the skilled artisan and they are included herein for purposes of illustration only. The purification protocol which is the subject of this invention is applicable, with only routine modification, to a variety of recombinant antibodies and antibody-like proteins regardless of how they are produced or cultured. For example a chimeric monoclonal antibody to CD4 was also
20 purified by the process of this invention.

The purified antibodies obtained by practicing the process of this invention have the following properties: 1) greater than 97% antibody protein by weight; 2) stable to proteolytic degradation at 4°C for at least three months; 3) low (< 0.1 E.U./mg protein) endotoxin; 4) low (< 1 pg/mg protein) DNA; 5)
25 non-antibody protein < 5% by weight; and 6) virally inactive. The following examples further illustrate this invention but are not offered by way of limitation of the claims herein.

EXAMPLE 1

INTRODUCTION

The procedure outlined below was developed for the isolation and purification of a monoclonal antibody against Respiratory Syncytial Virus (RSV).
5 This antibody is a "humanized" IgG expressed in CHO cells, and grown in a stirred tank bioreactor. The antibody is more fully described in PCT WO92/04381 and is otherwise referred to herein as RSHZ 19. The process is designed to prepare RSHZ-19 of >95% purity while removing contaminants derived from the host cell, cell culture medium, or other raw materials. The process in its most preferred
10 embodiment consists of three purification steps (Protein A affinity, cation exchange, and hydrophobic interaction chromatography), two viral inactivation steps, and a diafiltration step to exchange the product into a final buffer of choice (outlined in Figure 1). All steps are carried out at room temperature (18 - 25 °C). All buffers are prepared with WFI and filtered through either a 0.2 micron filter or a 10,000
15 MWCO membrane before use. Buffer formulations are listed in Table 1. Tables 2, 4, 6 and 8 show the column parameters for examples IA, IB, IC and ID respectively. Tables 3, 5 and 7 and 9 provide a purification summary for examples IA, IB, IC and ID respectively.

20 The first step in the process (Protein A affinity chromatography on ProSep A) can be rapidly cycled to accommodate varying amounts of cell-free culture fluid (CCF), and has a capacity of approximately 15 grams RSHZ-19 per liter of ProSep A. For example, 500 liters CCF containing 400 - 500 grams of IgG can be processed in 5 or 6 cycles. The downstream steps of the process (Cation Exchange
25 Chromatography (CEC) and Hydrophobic Interaction Chromatography (HIC) are scaled to accommodate approximately 130 - 140 grams RSHZ-19 per cycle. Thus, a 500 liter culture containing 400 - 500 grams of RSHZ-19 is processed in three downstream cycles after capture on ProSep A.

30 The hydrophobic interaction chromatography step (HIC) has been demonstrated to remove residual Protein A that leaches from the Protein A column during elution (See: examples IA-D). In addition, aggregates of IgG can be removed over HIC, as shown in examples IC and ID.

The process description is normalized for any scale; linear flow rates listed are independent of column diameter, loading ratios are in mass per unit column volume. Examples are provided for the operation and the recovery at 1 gram, 40 gram, and 125 gram scales. (Examples IA-ID).

5

Purification Process Description

Removal of Cells from Culture

10 To harvest the culture fluid, the cells are removed using a tangential-flow microfiltration device (Prostak) equipped with a 0.65 micron filter or equivalent. The product is recovered in the permeate. For small volume cultures, centrifugation can be used.

Affinity Capture by Protein A Chromatography

15 The IgG is recovered from the CCF by adsorption chromatography on a column of ProSep A (BioProcessing Ltd.) previously equilibrated with PBS. The medium is applied to the column at a flow rate up to 1000 cm/hr at a load ratio of up to 15 grams IgG per liter column volume. After loading the column, it is washed with at least 3 column volumes of PBS containing 0.1 M glycine. The RSHZ-19 is eluted with a low pH buffer by applying approximately 3 column volumes of Elution Buffer.

20 The Protein A chromatography removes a large proportion of cell and media derived impurities (particularly protein and DNA in the flow-through and wash fractions), and concentrates RSHZ-19 in the elution buffer for further processing.

Viral Inactivation at acid pH (Optional)

25 The Protein A column eluate is collected and adjusted to pH 3.5 by the addition of 2.5 M HCl. The solution is transferred to a second vessel and held at pH 3.5 for at least thirty minutes to provide viral inactivation, and readjusted to pH 5.5 by the addition of Tris buffer. The resulting solution is filtered through a prefilter (Millipore Polygard or equivalent) and a sterilized

30

35

0.2 µm filter (Millipore Millipak or equivalent), and held in sterile containers at 4 °C, or frozen and held at -70 °C.

5 The pH 3.5 treatment provides viral inactivation, and the pH 5.5 adjustment prepares the solution for cation exchange chromatography (CEC). The pH 3.5 treatment can be omitted if desired.

Cation Exchange Chromatography

10 The pH inactivated Protein A eluate is further purified by CEC chromatography on column of CM SEPHAROSE FF (Pharmacia LKB). The sample is applied to the equilibrated column at a flow rate of 150 cm/hr and a load ratio of ≤20 grams protein per liter CM SEPHAROSE. After loading, the column is washed with 3 to 5 column volumes of Equilibration Buffer.
15 The product is eluted with 3 - 5 column volumes of Elution Buffer.

The cation exchange chromatography step removes protein and non-protein impurities.

20

Viral Inactivation with Guanidine

The cation exchange eluate is adjusted to approximately 2.0 M guanidine hydrochloride by the slow addition (with mixing) of one-half volume of Guanidine Stock Solution. The rate of reagent addition is adjusted so that it
25 is added over a 5 - 15 minute period. The solution is transferred to a second vessel, and is held for thirty minutes to achieve viral inactivation. After holding, an equal volume of Ammonium Sulfate Stock Solution is slowly added (with mixing), and the hydrophobic interaction chromatography (HIC)
30 step is performed immediately. The rate of reagent addition is adjusted so that it is added over a 5 - 15 minute period.

The guanidine treatment provides a second viral inactivation step, when an acid inactivation step is employed, and keeps the RSHZ-19 soluble after ammonium sulfate addition; the addition of ammonium sulfate serves to
35 dilute the guanidine and prepare the solution for HIC.

Hydrophobic Interaction Chromatography

5 The guanidine-treated solution is further purified by application to an HIC column consisting of TOYOPEARL Phenyl-650M previously equilibrated with Equilibration Buffer. The guanidine-treated solution is applied to the column at a flow rate of 150 cm/hr and a load ratio of ≤ 20 grams protein per liter Phenyl-650M. After loading, the column is washed with 3 to 5 column volumes of Equilibration Buffer. A linear gradient of decreasing ammonium sulfate is applied at a flow rate of 100 - 150 cm/hr, and the RSHZ-19 elutes
10 as one major peak with impurities eluting later in the gradient. The slope of the gradient is approximately 20 column volumes, starting at 100% Equilibration Buffer and ending at 100% Gradient Buffer (1.3 to 0 M ammonium sulfate). The peak is collected until the absorbance decreases to 20% of the maximum peak absorbance, then collection of the product
15 fraction is ended. After the gradient ends, the column is washed with approximately 3 column volumes of Strip Buffer.

20 The HIC chromatography step removes additional protein and non-protein impurities, most notably residual Protein A, IgG aggregates, and host DNA.

Concentration, Diafiltration and Final Filtration

25 The HIC elute is concentrated to approximately 10 milligram per milliliter using a tangential-flow ultrafiltration device (such as a Millipore CUF) outfitted with a 30,000 molecular weight cut-off filter, diafiltered into a suitable formulation buffer and filtered through a sterilized 0.2 micron filter (Millipore Millipak or equivalent) into sterilized containers.

30 In-Process Assays

Process intermediates are assayed for total protein concentration by OD280 or Bradford assay, RSHZ-19 concentration by HLPC, sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). Aggregated
35 product is assessed by size exclusion HPLC on TSK3000 SWXL, and Protein A residue is assayed using an ELISA.

Pooling Criteria

5 The eluate fractions from the Protein A capture and cation exchange steps are pooled based on the UV tracing on the chromatogram, and the entire peak is collected. The eluate from the HIC step is pooled based on the UV tracing, and the main peak is pooled until the UV reading on the tailing side of the peak reaches 20% of the peak maximum. The HIC tail fraction contains the majority of the Protein A and aggregated IgG's.

Table 1: Buffer Formulations

	Buffer Name	Composition
5	PBS	20 mM sodium phosphate, 150 mM sodium chloride, pH 7
	PBS/glycine	PBS plus 0.1 M glycine
10	ProSep Elution Buffer	25 mM citrate, pH 3.5
	CM SEPHAROSE Equilibration Buffer ..	10 mM citrate, pH 5.5
15	CM SEPHAROSE Elution Buffer	40 mM citrate, 100 mM sodium chloride, pH 6
	Guanidine Stock Solution	6 M guanidine hydrochloride, 50 mM sodium phosphate, pH 7
20	2.6 M Ammonium Sulfate	2.6 M ammonium sulfate, 50 mM sodium phosphate, pH 7
	Stock Solution	
	2.0 M Ammonium Sulfate	2.0 M ammonium sulfate, 50 mM sodium phosphate, pH 7
	Stock Solution	
25	Phenyl-650 Equilibration Buffer	1.3 M ammonium sulfate, 50 mM sodium phosphate, pH 7
	Phenyl-650 Gradient Buffer	50 mM sodium phosphate, pH 7
30	Butyl-650 Equilibration Buffer	1.0 M ammonium sulfate, 50 mM sodium phosphate, pH 7
	Butyl-650 Gradient Buffer	50 mM sodium phosphate, pH 7
35	HIC Strip Buffer	0.2 M NaOH

Example IA. RSHZ-19 purification at 1 gram scale using TOYOPEARL Phenyl-650M

A 5.0 liter (20 cm diameter by 16 cm length) ProSep A affinity column was
5 equilibrated with PBS (see Table 1) at 5.2 liter/min. 100 liters of conditioned culture
medium containing 0.8 grams per liter of RSHZ-19 monoclonal antibody was
clarified by microfiltration as described above, and applied to the column at a flow
rate of 5.2 liter/min. After the load, approximately 15 liters of PBS/glycine was
applied to the column at the same flow rate. The IgG was eluted by applying 15 - 20
10 liters of ProSep A elution buffer. Fractions of the non-bound peak and the elution
peak were collected and assayed for IgG content using an HPLC assay. The eluate
was approximately 15 liters in volume, and contained approximately 5 milligrams
protein per milliliter.

15 Immediately after elution, the sample was adjusted to pH 3.5 by the addition of 2.5
M hydrochloric acid, held for approximately 30 minutes, and adjusted to pH 5.5 by
the addition of approximately 350 milliliters of 1 M Tris base. After neutralizing to
pH 5.5, the sample was filtered through a 0.1 micron Polygard CR filter in tandem
with a sterile 0.2 micron Millipak 200, into a sterile container. The filtrate was
20 stored at 4 °C. Samples of the filtrate were analyzed for IgG content using an HPLC
assay, and for total protein by absorbance at 280 nanometers. The samples were also
analyzed for Protein A content by an ELISA procedure. This pH 3.5 treated and
filtered Prosep A eluate was used as the CM SEPHAROSE load in Examples IA, B
and C.

25 400 milliliters of pH 3.5 treated and filtered ProSep A eluate were loaded directly
onto a 220 milliliter (4.4 cm diameter x 15 cm length) column of CM SEPHAROSE
FF at 38 mL/min, which had been previously equilibrated with CM Equilibration
buffer. After loading, the column was washed at 38 mL/min with approximately 700
30 milliliters of CM Equilibration Buffer. The IgG was eluted by applying CM Elution
Buffer at 38 mL/min. The IgG came off the column after approximately 1 bed
volume of Elution Buffer had passed. The entire peak was collected as CM
SEPHAROSE eluate. Fractions of the CM non-bound, eluate and strip fractions
were collected and analyzed for IgG content, total protein content, and Protein A
35 content as described previously. The eluate was approximately 160 milliliters in
volume, and contained approximately 12 milligrams protein per milliliter. This CM

SEPHAROSE eluate was split into two equal portions of approximately 80 milliliters, and was used in Examples IA and IB for the HIC load.

To 80 milliliters of CM SEPHAROSE eluate was added (slowly with constant stirring) a total of 40 milliliters of Guanidine Stock Solution. This brought the guanidine concentration to 2 M for viral inactivation. While stirring the guanidine-treated solution, a total of 120 milliliters of 2.6 M Ammonium Sulfate Stock Solution was added. The resulting solution was 1.0 M in guanidine and 1.3 M in ammonium sulfate. The ammonium sulfate treated solution was applied to an 80 mL column (3.2 cm diameter x 10 cm length) of TOYOPEARL Phenyl-650M, previously equilibrated with Phenyl Equilibration Buffer. The flow rate was 20 mL/min throughout the run. After loading, the column was washed with approximately 350 milliliters of Phenyl Equilibration Buffer. The IgG was eluted by applying a linear gradient starting at 85% Equilibration/15% Gradient buffer and ending at 0% Equilibration/100% Gradient buffer, in 18 - 19 column volumes. This represents a starting ammonium sulfate concentration of approximately 1.1 M and an ending concentration of 0 M. The slope of this gradient was approximately a 4.7% increase in elution buffer per column volume, or -0.061 M ammonium sulfate per column volume. The IgG began to elute from the column at approximately 7 column volumes and ended at approximately 13 column volumes into the gradient (ammonium sulfate concentration of approximately 0.7 to 0.3 M). The eluted fraction was collected until the UV absorbance on the tailing side of the peak decreased to 20% of the peak height, then collection was switched to another vessel (tail fraction). At the end of the gradient, approximately 250 mL of HIC Strip Buffer was applied to regenerate the column. Fractions of the Phenyl non-bound, eluate, tail and strip fractions were collected and analyzed for IgG content, total protein content, and Protein A content as described previously. The eluate was approximately 300 milliliters in volume, and contained approximately 2.4 milligrams protein per milliliter.

Table 2 summarizes the column parameters for this example. The product and protein recovery data for each step are shown in Table 3, along with the Protein A content, expressed as nanograms Protein A per milligram IgG (ng/mg). As seen in Table 3, the Protein A reduction over Phenyl-650M is approximately 4-fold, and the recovery is approximately 94%.

Table 2: Column Parameters at 1 gram scale using Phenyl-650M

5	Step	Column Volume (liter)	Column dia x length (cm)	Load Ratio	Flow Rates	
					(cm/hr)	(mL/min)
10	ProSep A	5.0	20 x 16	16.0 g IgG per liter bed volume	1000	5200
	CM SEPHAROSE FF	0.22	4.4 x 15	9.1 g protein per liter bed volume.	150	38
15	Phenyl-650M	0.08	3.2 x 10	10.4 g protein per liter bed volume	150	20

Table 3: Purification Summary for Example IA, 1 gram scale using Phenyl-650M

	Step	Volume (Liters)	Total RSHZ-19 ^a (Grams)	Total Protein ^b (Grams)	Step Yield (%)	Protein A ^c (ng/mg)
5	Cell-free Culture Fluid	100	80.3	n.d.	--	0
10	ProSep A Eluate	15.8	73.8	80.4	92	20.2
15	CM SEPHAROSE ^d (0.4) ^d Load		(1.87) ^d	(2.04) ^d		
	CM SEPHAROSE Eluate	0.16	2.01	1.88	100	14.5
20	Phenyl-650M ^d Load	(0.21) ^d	(0.72) ^d	(0.83) ^d		
	Phenyl-650M Eluate	0.31	0.68	0.73	94	3.5
25	(Phenyl Tail	0.59	0.075	0.10	--	133) ^e
	Cumulative Recovery (%)				86	
30	^a by HPLC ^b by Absorbance at 280 nm ÷ 1.27 mL mg ⁻¹ cm ⁻¹ ^c by ELISA ^d Only a portion of the total eluate from the previous column was carried forward, as described in the text above					
35	^e Protein A migrates primarily in the Tail fraction					

Example IB. RSHZ-19 purification at 1 gram scale using TOYOPEARL Butyl-650M

This preparation used the same CM SEPHAROSE eluate as described in Example IA, and the HIC step was performed using TOYOPEARL Butyl-650M instead of Phenyl-650M. The preparation of the CM SEPHAROSE eluate is described in Example IA above. To 80 milliliters of CM SEPHAROSE eluate was added (slowly with constant stirring) a total of 40 milliliters of Guanidine Stock Solution. This brought the guanidine concentration to 2 M for viral inactivation. While stirring the guanidine-treated solution, a total of 120 milliliters of 2.0 M Ammonium Sulfate Stock Solution was added. The resulting solution was 1.0 M in guanidine and 1.0 M in ammonium sulfate. The ammonium sulfate treated solution was applied to a an 80 mL column (3.2 cm diameter x 10 cm length) of TOYOPEARL Butyl-650M, previously equilibrated with Butyl Equilibration Buffer. The flow rate was 20 mL/min throughout the run. After loading, the column was washed with approximately 350 milliliters of Butyl Equilibration Buffer. The IgG was eluted by applying a linear gradient starting at 65% Equilibration/35% Gradient buffer and ending at 20% Equilibration/80% Gradient buffer, in 12 - 13 column volumes. This represents a starting ammonium sulfate concentration of approximately 0.65 M and an ending concentration of approximately 0.2 M. The slope of this gradient was approximately a 3.3% increase in elution buffer per column volume, or -0.033 M ammonium sulfate per column volume. The IgG began to elute from the column at approximately 2 column volumes and ended at approximately 9 column volumes into the gradient (ammonium sulfate concentration of approximately 0.58 to 0.35 M). The eluate fraction was collected until the UV absorbance on the tailing side of the peak decreased to 10% of the peak height, then collection was switched to another vessel. At the end of the gradient, approximately 250 mL of Butyl Gradient Buffer was applied, and a small peak eluted and was collected. Approximately 250 mL of HIC Strip Buffer was applied to regenerate the column. Fractions of the Butyl non-bound, eluate, tail and strip fractions were collected and analyzed for IgG content, total protein content, and Protein A content as described previously. The eluate was approximately 400 milliliters in volume, and contained approximately 1.5 milligrams protein per milliliter.

Table 4 summarizes the column parameters for this example. The product and protein recovery data for each step are shown in Table 5, along with the Protein A content, expressed as nanograms Protein A per milligram IgG (ng/mg). Although

the recovery of IgG is lower compared to Example IA (79% v. 94%), the Protein A content is reduced approximately 20-fold using Butyl-650M as an HIC step..

5

Table 4: Column Parameters at 1 gram scale using Butyl-650M

Step	Column Volume (liter)	Column dia x length (cm)	Load Ratio	Flow Rates	
				(cm/hr)	(mL/min)
10	CM SEPHAROSE FF	0.22	4.4 x 15	9.1 g protein per liter bed volume	150 38
15	Butyl-650M	0.08	3.2 x 10	10.4 g protein per liter bed volume	150 20

Table 5: Purification Summary for Example IB, 1 gram scale using Butyl-650M

	Step	Volume (Liters)	Total RSHZ-19 ^a (Grams)	Total Protein ^b (Grams)	Step Yield (%)	Protein A ^c (ng/mg)
5	Cell-free Culture Fluid	100	80.3	n.d.	--	0
10	ProSep A Eluate	15.8	73.8	80.4	92	20.2
	CM SEPHAROSE ^d (0.4) ^d Load		(1.87) ^d	(2.04) ^d		
15	CM SEPHAROSE Eluate	0.16	2.01	1.88	100	14.5
	Butyl-650M ^d Load	(0.21) ^d	(0.76) ^d	(0.86) ^d		
20	Butyl-650M Eluate	0.41	0.60	0.62	79	0.7
25	(Butyl Tail	0.40	0.03	0.03	--	31.4) ^{e,f}
	(Butyl Strip	0.53	0.10	0.10	--	n.d.) ^g
	Cumulative Recovery (%)				73	
30	Mass Balance (% of load)				95	
	^a by HPLC ^b by Absorbance at 280 nm ÷ 1.27 mL mg ⁻¹ cm ⁻¹ ^c by ELISA ^d Only a portion of the total eluate from the previous column was carried forward, as described above ^e Protein A migrates primarily in the Tail fraction ^f Tail contains 3% of the load ^g strip contains 13% of the protein that was loaded					
35						
40						

Example IC. RSHZ-19 purification at 40 gram scale using TOYOPEARL Phenyl-650M

This preparation used the same ProSep A eluate as described in Example IA, and the downstream steps were scaled-up to accommodate approximately 40 grams of protein, using TOYOPEARL Phenyl-650M as the HIC medium. The preparation of the CM SEPHAROSE Load is described in Example IA above. 7.8 liters of pH 3.5 treated and filtered ProSep A eluate were loaded directly onto a 4.2 liter (25 cm diameter x 8.5 cm length) column of CM SEPHAROSE FF which had been previously equilibrated with CM Equilibration buffer at 1.2 L/min. After loading, the column was washed at 1.2 L/min with approximately 8 liters of CM Equilibration Buffer. The IgG was eluted by applying CM Elution Buffer at 1.2 L/min. The IgG came off the column after approximately 1 bed volume of Elution Buffer had passed. The entire peak was collected as CM SEPHAROSE eluate. The column fractions of the CM non-bound and eluate were collected and analyzed for IgG content, total protein content, and Protein A content as described previously. The eluate was approximately 5.7 liters in volume, and contained approximately 6 - 7 milligrams protein per milliliter.

To 5.6 liters of CM SEPHAROSE eluate was added (slowly with constant stirring) a total of 2.8 liters of Guanidine Stock Solution (3.2 kilogram by weight). This brought the guanidine concentration to 2 M for viral inactivation, at a volume of 8.3 liters. While stirring the guanidine-treated solution, a total of 8.3 liters (9.7 kg by weight) of 2.6 M Ammonium Sulfate Stock Solution was added. The resulting solution was 1.0 M in guanidine and 1.3 M in ammonium sulfate, with a final volume of 16.7 liters. The ammonium sulfate treated solution was applied to a 4.6 liter column (18 cm diameter x 18 cm length) of TOYOPEARL Phenyl-650M, previously equilibrated with Phenyl Equilibration Buffer. The flow rate was 0.5 - 0.6 L/min throughout the run. After loading, the column was washed with approximately 14 liters of Phenyl Equilibration Buffer. The IgG was eluted by applying a linear gradient starting at 100% Equilibration Buffer and ending at 100% Gradient buffer, in 20 column volumes. This represents a starting ammonium sulfate concentration of approximately 1.3 M and an ending concentration of 0 M. The slope of this gradient was approximately a 5% increase in elution buffer per column volume, or -0.065 M ammonium sulfate per column volume. The IgG began eluting from the column at approximately 7 column volumes and ended at approximately 12

column volumes into the gradient (ammonium sulfate concentration of approximately 0.85 to 0.5 M). The eluate fraction was collected until the UV absorbance on the tailing side of the peak decreased to 20% of the peak height, then collection was switched to another vessel (tail). Fractions of the Phenyl non-bound, eluate and tail and strip fractions were collected and analyzed for IgG content, total protein content, and Protein A content as described previously. The eluate was approximately 15.4 L in volume, and contained approximately 2.2 milligrams protein per milliliter.

The Phenyl Eluate was concentrated to approximately 16 mg/mL using a tangential flow ultrafiltration apparatus (CUF, Millipore Corp.) equipped with 30,000 MWCO Omega membranes (Filtron Corp.) and buffer exchanged by continuous diafiltration against a suitable formulation buffer.

Table 6 summarizes the column parameters for this example. The product and protein recovery data for each step are shown in Table 7, along with the Protein A content, expressed as nanograms Protein A per milligram IgG (ng/mg) and the IgG aggregate content, expressed as % of total IgG. As seen in Table 7, the Protein A reduction over Phenyl-650M is approximately 3-fold, and the recovery is approximately 90%. IgG aggregates were reduced from 0.5% in the CM SEPHAROSE eluate to 0.06% in the formulated product.

Table 6: Column Parameters at 40 gram scale

Step	Column Volume (liter)	Column dia x length (cm)	Load Ratio	Flow Rates	
				(cm/hr)	(L/min)
CM SEPHAROSE FF	4.2	25 x 8.5	8.9 g protein per liter bed volume	150	1.2
Phenyl-650M	4.6	18 x 18	10.4 g protein per liter bed volume	140	0.6

Table 7: Purification Summary for Example IC: 40 gram scale

	Step	Volume (Liters)	Total RSHZ-19 ^a (Grams)	Total Protein ^b (Grams)	Step Yield (%)	Protein A ^c (ng/mg)	IgG Aggregate (%)
5	Cell-free Culture Fluid	100	80.3	n.d.	--	0	n.d.
10	ProSep A Eluate	15.8	73.8	80.4	92	20.2	n.d.
	CM SEPHAROSE ^d (7.84) ^d Load		(36.0) ^d	(37.3) ^d			
15	CM SEPHAROSE Eluate	5.71	36.5	37.3	100	21.4	0.5%
20	Phenyl-650M Eluate (Product)	15.4	32.8	33.6	90	8.0	<0.05
	Phenyl Tail	24.6	2.5	3.0	--	78.0	5.1) ^e
25	Formulated Product	2.0	32.8	32.0	100	6.5	0.06
	Cumulative Recovery (%)				83		
	Mass Balance (% of load)				97		
30	-----						
	^a by HPLC						
	^b by Absorbance at 280 nm ÷ 1.27 mL mg ⁻¹ cm ⁻¹						
	^c by ELISA						
	^d Only a portion of the total ProSep A Eluate was carried forward						
35	^e Protein A and IgG aggregates migrate primarily in the tail fraction						

Example ID. RSHZ-19 purification at 125 gram scale using TOYOPEARL Phenyl-650M

A 5.5 liter (20 cm diameter by 18 cm length) ProSep A affinity column was equilibrated with PBS (see Table 1) at 4.8 liter/min. 450 liters of conditioned culture medium containing 0.94 grams per liter of RSHZ-19 monoclonal antibody was clarified by microfiltration as described above, and applied in four separate 90 - 95 liter portions and one 40 liter portion to the column at a flow rate of 4.8 liter/min (and so throughout). Each cycle on the column ran as follows: After the load, approximately 17 liters of PBS/glycine was applied to the column at the same flow rate. The IgG was eluted by applying 15 - 20 liters of ProSep A Elution buffer. Fractions of the non-bound peak and the elution peak were collected and assayed for IgG content using an HPLC assay. The eluate from each cycle was approximately 9 liters in volume, and contained approximately 5 - 10 milligrams protein per milliliter. Immediately after elution, the ProSep A eluates were adjusted to pH 3.5 by the addition of 2.5 M hydrochloric acid, held for approximately 30 minutes, and adjusted to pH 5.5 by the addition of approximately 250 milliliters of 1 M Tris base. After neutralizing to pH 5.5, the eluates were pooled together, and filtered through a 0.1 micron Polygard CR filter in tandem with a sterile 0.2 micron Millipak 200, into 5 liter aliquots in sterile containers. The filtrate was stored at 4 °C. Samples of the filtrate were analyzed for IgG content using an HPLC assay, and for total protein by absorbance at 280 nanometers. The samples were also analyzed for Protein A content by an ELISA procedure, and IgG aggregates by HPLC.

The downstream steps were scaled-up to accommodate approximately 120 - 140 grams of protein. 16.3 liters of pH 3.5 treated and filtered ProSep A eluate containing approximately 130 grams of protein was loaded directly onto a 14.4 liter (35 cm diameter x 15 cm length) column of CM SEPHAROSE FF at 2.4 L/min, which had been previously equilibrated with CM Equilibration buffer. After loading, the column was washed at 2.4 L/min with approximately 45 liters of CM Equilibration Buffer. The IgG was eluted by applying CM Elution Buffer at 2.4 L/min. The IgG began to elute from the column after approximately 1 - 2 bed volumes of Elution Buffer had passed. The entire peak was collected as CM SEPHAROSE eluate. Fractions of the CM non-bound and eluate were collected and analyzed for IgG content, total protein content, and IgG aggregate. The eluate was approximately 21 liters in volume, and contained approximately 120 grams protein.

To 19.4 liters of CM SEPHAROSE eluate was added (slowly with constant stirring) a total of 9.7 liters of Guanidine Stock Solution. This brought the guanidine concentration to 2 M for viral inactivation, at a volume of 29.1 liters. While stirring the guanidine-treated solution, a total of 29.1 liters of 2.6 M Ammonium Sulfate Stock Solution was added. The resulting solution was 1.0 M in guanidine and 1.3 M in ammonium sulfate, with a final volume of 58.2 liters. The ammonium sulfate treated solution was applied to a 12.4 liter column (30 cm diameter x 18 cm length) of TOYOPEARL Phenyl-650M, previously equilibrated with Phenyl Equilibration Buffer. The flow rate was 1.1 - 1.3 L/min throughout the run. After loading, the column was washed with approximately 37 liters of Phenyl Equilibration Buffer. The IgG was eluted by applying a linear gradient starting at 80% Equilibration Buffer/20% Gradient Buffer and ending at 20% Equilibration Buffer/80% Gradient buffer, in 12 column volumes. This represents a starting ammonium sulfate concentration of approximately 1.0 M and an ending concentration of approximately 0.26 M. The slope of this gradient was approximately a 5% increase in Gradient buffer per column volume, or -0.065 M ammonium sulfate per column volume. The IgG came off the column essentially in the middle of the gradient, with the peak containing approximately 0.8 M ammonium sulfate. The eluate fraction was collected until the UV absorbance on the tailing side of the peak decreased to 20% of the peak height, then collection was switched to another vessel (tail). Fractions of the Phenyl non-bound, eluate and tail and strip fractions were collected and analyzed for IgG content, total protein content, and IgG aggregate. The eluate was approximately 29 L in volume, and contained approximately 100 grams protein.

The Phenyl Eluate was concentrated to approximately 10 mg/mL using a tangential flow ultrafiltration apparatus (CUF, Millipore Corp.) equipped with 30,000 MWCO Omega membranes (Filtron Corp.) and buffer exchanged by continuous diafiltration against a suitable formulation buffer. The product was analyzed for IgG content, total protein, Protein A and IgG aggregate.

Table 8 summarizes the column parameters for this example. The product and protein recovery data for each step are shown in Table 9, along with the Protein A content, expressed as nanograms Protein A per milligram IgG (ng/mg) and the IgG aggregate content, expressed as % of total IgG. As seen in Table 9, the Protein A reduction over CM SEPHAROSE and Phenyl-650M is approximately 7-fold, and

the cumulative recovery is approximately 70%. IgG aggregates were reduced from 0.4% in the CM SEPHAROSE eluate to 0.06% in the formulated product.

5

Table 8: Column Parameters at 125 gram scale

Step	Column Volume (liter)	Column dia x length (cm)	Load Ratio	Flow Rates	
				(cm/hr)	(L/min)
ProSep A	5.0	20 x 18	14 - 16 g RSHZ-19 per liter bed volume	915	4.8
CM SEPHAROSE FF	14.4	35 x 15	8.4 g protein per liter bed volume	150	2.4
Phenyl-650M	12.4	30 x 18	8.6 g protein per liter bed volume	100	1.2

20

Table 9: Purification Summary for Example ID: 125 gram scale

	Step	Volume (Liters)	Total RSHZ-19 ^a (Grams)	Total Protein ^b (Grams)	Step Yield (%)	Protein A ^c (ng/mg)	IgG Aggregate (%)
5	Cell-free Culture Fluid	416	392	477 ^c	--	0	n.d.
10	ProSep A Eluate	48.7	375	384	96	11.7	0.4
	CM SEPHAROSE ^d (16.3) ^d Load		(125) ^d	(129) ^d	--		
15	CM SEPHAROSE Eluate	20.6	116	117	93	n.d.	0.4
20	Phenyl-650M Eluate	29.1	98.1	99.8	85	n.d.	<0.05
	Formulated Product	9.29	89.8	91.1	92	1.7	0.06
25	Cumulative Recovery (%)				70		

^a by Reversed-Phase HPLC^b by Absorbance at 280 nm ÷ 1.27 mL mg⁻¹ cm⁻¹^c by Bradford assay

30 ^d Downstream process capacity is approximately 140 grams; only a portion of the total ProSep A eluate is carried forward

Table 10: Purity analysis: 125 Gram Scale

5	Step	Aggregates	Purity ^a	Activity ^b
		(% of Total IgG)	(% of Total Area)	(%)
	CCF	not applicable	not done	80 ^c
	ProSep Eluate	0.4	98.5	105
10	CM Eluate	0.4	98.4	109
	Phenyl Eluate	<0.05	98.3	99
15	Final product	0.06	99.7	115

^a Determined by scanning densitometry of reducing SDS-PAGE; sum of area of Heavy and Light chains of IgG

^b Calculated ratio of activity (determined by Bovine RS Virus binding ELISA) to RSHZ-19 concentration (determined by A280)

^c Calculated ratio of activity, by bovine RS virus ELISA, to RSHZ-19 concentration by HPLC

EXAMPLE II

The anti-CD4 monoclonal antibody CH-CD14 was made by cell culture techniques and partially purified using Protein A and ion exchange chromatography in a fashion similar to that used in Example I, except that an anion exchange resin was employed. An Amicon column (1 cm diameter by 10 cm high) was packed with 8 mL of Phenyl TOYOPEARL 650M resin (lot # 65PHM01 H). The flow rate was maintained at 2 mL/min for all steps. The column was equilibrated with 20 mL Equilibration buffer (1 M ammonium sulfate, 50 mM sodium phosphate, pH 7.0). The product was prepared for loading on the column by taking 5 mL partially purified CH-CD4 monoclonal antibody (17 mg/mL concentration by absorbance at 280 nm), adding 2.5 mL 6 M guanidine HCl, 50 mM sodium phosphate, pH 7.0, mixing, holding for 30 minutes, and then adding 7.5 mL 2 M ammonium sulfate, 50 mM sodium phosphate, pH 7.0. The final ammonium sulfate concentration of the load was 1 M, the final guanidine HCl concentration was 1 M, the final volume after samples were taken was 12 mL, and the final concentration of product was 5.6 mg/mL. This material was then loaded on the column at 2 mL/min and then eluted with a 10 column volume linear gradient of Equilibration buffer to 50 mM sodium phosphate, pH 7.0. Fractions of approximately 4 mL were taken as the product eluted from the column.

The results are shown in Table 11. All of the product eventually eluted in the gradient. Protein A also eluted and was enriched in later fractions from the gradient. To achieve a reduction in Protein A in the final product, it was necessary to exclude some of the fractions at the end of the gradient, thus reducing the yield of product. A two-fold reduction in Protein A was possible with an 86% yield of product and a three-fold reduction was possible with a 50% yield.

TABLE 11

Fraction N .	Volume (mL)	Product (mg/mL)	Protein A (ng/mL)	Cumulative Specific Protein A (ng/mg)	Cumulative Product Yield	Protein A Removal Factor ^a
	Load					
0	12	5.8	200.7	35		
	Eluate					
1	5.5	0.04	0.0	0	0%	
2	3	0.31	0.0	0	2%	
3	4.2	1.32	6.4	4	10%	8.6
4	4.2	3.1	16.9	5	28%	7.0
5	4	4	68.3	10	51%	3.3
6	4.2	3.4	80.9	14	72%	2.4
7	4.2	2.3	94.1	19	86%	1.9
8	4.2	1.3	79.3	22	94%	1.6
9	4.2	0.71	49.9	24	98%	1.4
10	4.2	0.36	32.8	26	100%	1.3
11	4	0.19	22.9	27	101%	1.3
12	4	0.11	13.5	27	102%	1.3
13	9.6	0.05	8.5	28	102%	1.2

- ^a Removal factor is calculated by pooling eluate fractions from start of eluate to desired fraction and dividing initial Protein A in load by Protein A ng/mg in pooled eluate fractions. For example, the factor 8.6 is calculated by dividing the sum of Protein A in fractions 1-3 by the sum of product in fractions 1-3 to get 4 ng/mg. This number is then divided by 35 ng/mg in the load to obtain 8.6.

EXAMPLE IIB:

CH-CD4 monoclonal antibody was partially purified, and prepared for loading on the HIC column as described in example IIA. The same column, flowrate, equilibration and loading described in example IIA were used. In this example, after the column was loaded and washed with Equilibration buffer, it was washed with 18 mL Wash buffer (1 M ammonium sulfate, 50 mM sodium citrate, pH 3.5). This was followed by another washing with 13.5 mL Equilibration buffer (described in example IIA). The column was eluted with a gradient and then washed with water as described in example IIA. The column eluate was divided into 14 mL fractions which were then analyzed for product and Protein A.

The results are presented in Table 12. Protein A could be reduced by 6 fold at 90% yield, and by 8 fold at 78% yield.

TABLE 12

Fraction No.	Volume (mL)	Product (mg/mL)	Protein A (ng/mL)	Cumulative Specific Protein A (ng/mg)	Cumulative Product Yield	Protein A Removal Factor ¹
	Load					
0	12	5.80	167.1	29		
	Eluate					
1	14	1.50	0	0.0	30%	—
2	14	2.35	13.6	3.5	78%	8.1
3	14	0.38	8.3	5.2	85%	5.6
4	14	0.18	0	5.0	89%	5.8
5	9	0.13	0	4.9	91%	5.9

¹ As in footnote "a" to Table 11.

Comparing the results of example IIB with example IIA it can be seen that Protein A was reduced by 6 to 8 fold with about 80% yield when the pH 3.5 wash was included, but reduction was only about 2 fold with 80% yield of CH-CDH without the pH 3.5 wash.

5 EXAMPLE IIC:

This example was performed similar to example IIB, except that the scale was increased. Equilibration and Wash buffers are described in example IIB. The column was 5 cm in diameter and 28 cm high and the flowrate was 50 mL/min. CH-CD4 monoclonal antibody was prepared and partially purified, as described in
10 example IIA. The partially purified product (440 mL) was mixed with 220 mL 6 M guanidine HCl for 31 min. Then, 660 mL 2 M ammonium sulfate, 50 mM sodium phosphate, pH 7.0 was added. The final ammonium sulfate concentration was 1 M and the final antibody concentration was 5.3 mg/mL. After sampling, the load volume was 1,290 mL.

15 The column was equilibrated with 2 column volumes of Equilibration buffer and then loaded on the column. The column was then washed with 630 mL of Equilibration buffer, 1,000 mL of Wash buffer, 800 mL of Equilibration buffer, and then eluted with a 5 column volume gradient from 0.75 M ammonium sulfate, 50 mM sodium phosphate, pH 7.0, to 50 mM sodium phosphate, pH 7.0. Fractions
20 were collected during elution.

The results are presented in Table 13. Protein A was reduced 100 fold with a yield of 70%, or by 30 fold with an antibody yield of 80%.

TABLE 13

Fraction No.	Volume (mL)	Product (mg/mL)	Protein A (ng/mL)	Cumulative Specific Protein A (ng/mg)	Cumulative Product Yield	Protein A Removal Factor
	Load					
0	1290	5.30	198	37		
	Eluate					
1	581	0.03	0	0.0	0%	
2	304	3.60	0	0.0	16%	
3	243	5.40	0	0.0	35%	
4	238	4.50	0	0.0	51%	
5	237	3.40	2.4	0.1	63%	282
6	239	2.2	4.7	0.4	71%	107
7	256	1.2	4.8	0.6	75%	66
8	252	0.7	7.6	0.9	78%	41
9	240	0.4	5.3	1.1	79%	33
10	250	0.4	5.7	1.4	81%	27
11	202	0.4	4.38	1.5	82%	25
12	210	1	148.5	6.8	85%	5

EXAMPLE IID

Partially purified CH-CD4 monoclonal antibody was prepared as shown in example IIA. Equilibration and Wash buffers are described in example IIB. Four milliliters of 6 M guanidine HCl, 50 mM sodium phosphate, pH 7.0 was added to 8
5 mL of a 9.5 mg/mL solution of partially purified CH-CD4 monoclonal antibody and incubated for 30 minutes. Then, 12 mL of 2 M ammonium sulfate, 50 mM sodium phosphate, pH 7.0 was added slowly. The column was 0.5 cm in diameter and 20 cm high (4 mL) and the flowrate for all steps was 0.5 mL/min. The column was rinsed with 2 column volumes each of water and Equilibration buffer. Then, 22 mL
10 of the load solution was passed through the column, followed by 2 column volumes of Equilibration buffer, followed by Wash buffer until the pH of the column effluent was 3.5. This was followed by Equilibration buffer until the effluent pH was 7.0. The column was eluted with 0.3 M ammonium sulfate, 50 mM sodium phosphate, pH 7.0. After the UV trace started to rise, 12.6 mL of eluate were collected and
15 analyzed for product and Protein A. The yield of product was 80% and the Protein A was reduced from 28 ng/mg to 6 ng/mg, a reduction of 4.7 fold.

WHAT IS CLAIMED IS:

1. A method for purifying monomeric IgG antibody from a mixture containing same comprising contacting said mixture with a hydrophobic interaction chromatographic support and selectively eluting the monomer from the support.
2. The method according to Claim 1 wherein the IgG is selected from the group consisting of anti-RSHZ-19 and CH-CD4.
3. The method according to Claim 1 wherein the HIC support is selected from the group consisting of alkyl_{C2-C8} agarose, aryl-agarose, alkyl-silica, aryl-silica alkyl organic polymer resin and aryl organic polymer resin.
4. The method according to Claim 3 wherein the support is selected from the group consisting of butyl-, phenyl-, and octyl-agarose and butyl-, phenyl- and ether- organic polymer resin.
5. The method according to Claim 4 wherein the support is phenyl-organic polymer resin.
6. The method according to Claim 4 wherein the support is butyl-organic polymer resin.
7. The method according to Claim 1 wherein the antibody is selectively eluted with a low salt buffer.
8. The method according to Claim 7 wherein the antibody is selectively eluted with a gradient decreasing in salt to 50 mM phosphate, pH7.0.
9. A method for the purification of an IgG antibody from conditioned cell culture medium containing same comprising sequentially subjecting the medium to (a) Protein A affinity chromatography, (b) ion exchange chromatography, and (c) hydrophobic interaction chromatography.
10. The method according to Claim 9 wherein the ion exchange chromatography employs a support selected from the group consisting of CM-23-, CM-32-, CM-52-cellulose; CM-and, SP-cross-linked dextrans, CM- and S-argose; CM-organic polymer resin; DEAE-QAE-Q-cross-linked dextians; DEAE-,QAE-,Q- linked agarose; and DEAE organic polymer resins and is by a buffered salt solution.
11. The method according to Claim 10 wherein the support is CM-agarose Fast Flow and the salt is NaCl.
12. The method according to Claim 10 wherein the buffered salt solution is 40 mM citrate containing, 100mM, NaCl, pH 6.0.

13. The method according to Claim 9 wherein the hydrophobic interaction chromatographic employs a support selected from the group consisting of alkyl C2-C8-agarose, aryl-agarose, alkyl-silica, aryl-silica, alkyl-organic polymer resin and aryl-organic polymer resin.

5 14. The method according to Claim 13 wherein the support is selected from the group consisting of butyl-, phenyl- and octyl-agarose and butyl-, phenyl- and ether-organic polymer resin.

15. The method according to Claim 14 wherein the support is phenyl-organic polymer resin or butyl-organic polymer resin.

10 16. The method according to Claim 9 wherein the support is phenyl- or butyl-organic polymer resin and the antibody is selectively eluted with a low salt buffer.

15 17. The method according to Claim 16 wherein the antibody is selectively eluted with a gradient decreasing to 50 mM sodium phosphate buffer, pH 7.0.

18. The method according to Claim 9 wherein the Protein A chromatography employs as a support Protein A linked to controlled pore glass and elution is by a low pH buffer.

20 19. The method according to Claim 18 wherein said buffer is 25 mM citrate, pH 3.5.

20. A method for purifying antibody from a conditioned cell medium comprising:

- (a) adsorbing the antibody onto a Protein A chromatographic support;
- (b) washing the adsorbed antibody with at least one buffer;
- 25 (c) eluting the antibody from step (b);
- (d) adsorbing the antibody from step (c) onto an ion exchange chromatographic support;
- (e) washing the adsorbed antibody with at least one buffer;
- (f) selectively eluting the antibody from step (e);
- 30 (g) adsorbing the eluate of step (f) onto a hydrophobic interaction chromatographic support;
- (h) washing the adsorbed antibody with at least one buffer;
- (i) eluting the adsorbed antibody; and
- (j) recovering the antibody.

35 21. The method according to Claim 20 which includes one or more optional steps of inactivating viruses if present.

22. The method according to Claim 21 wherein a viral inactivation step is performed after step (f) and before step (g).

23. The method according to Claim 22 wherein said viral inactivation step comprises treatment of the eluate with guanidine hydrochloride for a period
5 of time sufficient to inactivate virus followed by the addition of an ammonium sulfate solution.

24. The method according to Claim 23 wherein the guanidine hydrochloride is present at 2.0 M and following treatment eluate from step (f) is adjusted to 1.3 M ammonium sulfate.

10 25. The method according to Claim 22 wherein an additional viral inactivation step is performed after step (c) and before step (d).

26. The method according to Claim 25 wherein said additional viral inactivation step comprises treatment of the eluate of step (c) with acid.

27. The method according to Claim 26 wherein the pH of the eluate
15 is adjusted to pH 3.5 and maintained at that pH for a period of time sufficient to inactivate virus, and terminating the treatment by adjusting the pH to 5.5.

28. The method according to Claim 20 wherein the ion exchange support of step (d) is selected from the group consisting of carboxymethyl (CM), sulfoethyl (SE), sulfopropyl (SP), phosphate(P), diethylaminoethyl (DEAE),
20 quaternary aminoethyl (QAE), and quaternary (Q), substituted cellulosic resins, cross linked dextrans, agarose and organic polymer resins.

29. The method according to Claim 28 wherein the cationic support is CM-agarose.

30. The method according to Claim 20 wherein the hydrophobic
25 interaction chromatographic support is selected from the group consisting of alkyl C2-C8-agarose, aryl-agarose, alkyl-silica, aryl-silica, alkyl-organic polymer resin and aryl-organic polymer resins.

31. The method according to Claim 30 wherein the support is selected from the group consisting of butyl-, phenyl- and octyl-agarose and
30 phenyl-, ether- and butyl-organic polymer resins.

32. The method according to Claim 31 wherein the support is phenyl- or butyl-organic polymer resins

33. The method according to Claim 20 wherein said protein is recovered by pooling and concentrating the protein containing fractions from
35 chromatography step (i) by ultrafiltration.

34. The method according to Claim 20 wherein the chromatographic support of step (a) is Protein A linked to controlled pore glass.

35. The method according to Claim 20 wherein the absorbed antibody of step (h) is washed with two buffers, a first equilibration buffer and a second low pH wash buffer.

36. The method according to Claim 35 wherein the pH of the second
5 buffer is less than 4.0.


37. The method according to Claim 36 wherein the second buffer is 1 M ammonium sulfate, 50 mM sodium citrate, pH 3.5.

38. A method for removing Protein A from a mixture comprising Protein A and antibodies comprising contacting said mixture with a hydrophobic
10 interaction chromatography support and selectively eluting the antibody from the support.

39. The method according to Claim 38 which includes washing the support prior to elution with a buffer having a pH less than 7.0.

40. The method according to Claim 39 wherein the pH of the wash
15 buffer is less than 4.0.

41. The method according to Claim 40 wherein the buffer is (1 M ammonium sulfate, 50 mM sodium citrate, pH 3.5).

Conditioned Media**FIGURE 1**

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/01823

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :B01D 15/08

US CL : 210/635, 656; 530/390.5, 413, 417

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 210/635, 656, 198.2; 530/387.1, 390.5, 413, 417

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 5,164,487 (KOTHE ET AL.) 17 November 1992, column 3, lines 64-66	1-41
Y	US, A, 5,252,216 (FOLENA-WASSERMAN ET AL.) 12 October 1993, column 2, lines 53-59	1-41



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A		document defining the general state of the art which is not considered to be part of particular relevance
* E	* X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* L	* Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* O		document referring to an oral disclosure, use, exhibition or other means
* P	* &	document published prior to the international filing date but later than the priority date claimed

Date of the actual completion of the international search

23 MARCH 1995

Date of mailing of the international search report

11 APR 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ERNEST G. THERKORN

Telephone No. (703) 308-0962